

PBDEs – Rising Levels in Fish, Tox Review and the California Ban

U.S. EPA National Forum on
Contaminants in Fish
San Diego, California
January 28, 2004

Tom A. McDonald, M.P.H., Ph.D.
OEHHHA, California EPA
tmcdonal@oehha.ca.gov




Office of Environmental Health Hazard Assessment



1

The figure that started it all for us in California . . .


Organohalogen Compounds in Human Milk in Sweden
(Norén and Mieronyté, 1998)



Introduction



The polybrominated dipenylethers (PBDEs)



- Added to many consumer products
- Flame retardant
- Saves lives



Office of Environmental Health Hazard Assessment



PBDE Use in the Americas


PBDE Technical Mixture	Million lbs/yr (2001)	% of World's Use	Product Uses
Penta-BDE [†]	15.7	95	Furniture (foam cushions)
Octa-BDE	3.3	40	Electronics (ABS plastic, cable)
Deca-BDE	54.0	44	Electronics (HIPS plastic) and textiles

[†] Highly bioaccumulative


BSEF, 2003




Structural Similarity of PBDEs, Their Metabolites and Environmental Derivatives to T4 and PCBs




2,2',4,4',5,5'-hexachlorobiphenyl
(PCB-153)




Thyroxine (T4)




2,2',4,4',6-pentabromodiphenylether
(PBDE-100)



2-(2',4'-dibromophenoxy)-4,6-dibromoanisole
(methoxy-PBDE)



4-(2',4',6'-tribromophenoxy)-2,6-dibromophenol
(hydroxy-PBDE)



2,3,4,7,8-pentabromodibenzofuran
(PBDF)


5

Office of Environmental Health Hazard Assessment



The data that accelerated our efforts in CA

Lipid-normalized PBDE-47 (tetra) in Human Tissues



CA data from She *et al.* (2002)

6

Office of Environmental Health Hazard Assessment



PBDEs Relate to Several Important Topics of the Day

- **Children's Health**
- **Endocrine Disruption**
- **Persistent Organic Pollutants (POP)/ Persistent Bioaccumulative Toxicants (PBT)**
- **Emerging Environmental Challenge**
- **High Production Volume (HPV) Chemical**

7

Office of Environmental Health Hazard Assessment



PBDEs Have Become Ubiquitous Environment Contaminants

PBDEs are measured in


- Indoor and outdoor air
- Remote Arctic regions (i.e., long-range transport)
- House and office dust
- Rivers and lakes and sediments
- Sewage sludge
- Foods
- Biota (terrestrial and marine mammals, fish, humans)

8

Office of Environmental Health Hazard Assessment



Time-trend: PBDEs in Blubber of California Seals (She *et al.*, 2002)



N=11


9

Office of Environmental Health Hazard Assessment



Fish from San Francisco Bay

PBDE levels in striped bass and halibut 1997 and 2002




10

EWG (2003)

Office of Environmental Health Hazard Assessment



PBDEs in Columbia River Whitefish




11

Office of Environmental Health Hazard Assessment

Rayne *et al.* (2003) Environ Sci Technol 37(13):2847-54.



PBDEs in Lake Ontario Trout (1978 - 1998)




Luross *et al.* (2000)

12

Office of Environmental Health Hazard Assessment




PBDEs in Herring Gull Eggs - Great Lakes



13


PBDE Levels Are Rising in U.S. Residents (Sum 7 PBDE Congeners in Serum)



Sjodin *et al.* (2003)

Office of Environmental Health Hazard Assessment

14



High-end Human Exposures

- 5% of people likely have lipid-normalized PBDE levels greater than 300 ng/g
 - That's about 15 million people in the U.S.
- WHY?
 - Fish intake?
 - Indoor exposures, house dust?
 - Differences in pharmacokinetics (i.e., inter-individual variability in uptake, metabolism or excretion)?
 - Look for future research to address this question



For Many Individuals, PBDE Tissue Levels Have Now Surpassed PCB Levels

- **Initial data from our agency indicate that among 57 California women :**

**7% had higher tissue concentrations
of total PBDEs than total PCBs**



Toxicity Concerns for the PBDEs

- **Endocrine disruption**
 - Thyroid and estrogenic effects
- **Developmental effects**
 - Brain and reproductive organs
- **Possibly cancer**
 - NTP initiating long-term studies of Penta
 - Environmental conversion to dioxins/furans
 - Brominated dioxins/furans measured in people



Thyroid Hormone Disruption

- Good evidence in rats and mice
- Some evidence in humans
- Relative potencies
 - penta-BDE > octa-BDE >>> deca-BDE
- Effects additive with co-exposures to PCBs
- Possible mechanisms
 - Hormone mimicry (transthyretin binding)
 - UDGPT enzyme induction (\uparrow T4 excretion)


19

Office of Environmental Health Hazard Assessment



Example: Thyroid Hormone Disruption

- Zhou et al. (2002): Penta-BDE (tech.) given to pregnant rats 0, 1, 10, 30 mg/kg-d from gestational day 6 through postnatal day 21



Office of Environmental Health Hazard Assessment



Estrogenic Effects


- **Postnatal exposure of rats to PBDE-99 altered expression of estrogen-regulated genes** (Lichtensteiger *et al.*, 2003)
 - Prostate: androgen receptor, estrogen receptor ER- α and ER- β , insulin-like growth factor (IGF-I)
 - Brain: Progesterone receptor, ER- α
- ***In vivo* estrogenic activity was not predicted from *in vitro* assays.**
 - PBDE-99 low estrogenic activity in MCF-7 cells.



Developmental Toxicity

- **Neurological system (3 independent laboratories)**
 - Altered behavior, learning and memory in mice
 - Hearing loss in rats
 - Effects permanent, i.e., measured in adulthood
 - Effects additive with co-exposure of PBDE and PCB
- **Male reproductive system (2 labs):**
 - Delayed puberty
 - Increased ventral prostate and seminal vesicle weights
- **Female reproductive system (3 labs):**
 - Delayed puberty
 - Alterations to ovary cell structure





Risk:

Compare High-end Human Levels to Tissue Levels in PBDE-treated Rodents

- Estimates of rodent body burdens of PBDE resulting from doses that caused these effects are equivalent to total PBDE levels attained in humans.
 - If humans are as sensitive as animals, then there is:

“No margin of safety.” Science News (2003) 164:266-8.
 - Better data are needed to compare tissue concentrations between rodents and humans.

Office of Environmental Health Hazard Assessment

Risk (continued)

- **An even greater concern: PBDEs and PCBs may be working together.**
 - PCB levels are usually higher than PBDEs
 - Same effects on some mechanistic endpoints
 - Co-administration of PCB and PBDE caused additive effects with respect to:
 - **behavior alterations in mice**
 - **thyroid hormone disruption**
 - **PBDEs/PCB co-exposures further increases the likelihood that exposure will result in health effects.**

25

Office of Environmental Health Hazard Assessment



Penta- and Octa-BDE are Now Banned Chemicals



- **Banned in California starting 2008**
 - AB302 (Chan et al.), signed into law in August.
- **U.S. manufacturer announced it will voluntarily cease production by end of 2004**
- **Banned by the European Union starting 2005**
- **Already voluntarily phased out in Japan**



Office of Environmental Health Hazard Assessment



Text of California Ban (AB302)

“ On and after January 1, 2008, a person may not manufacture, process, or distribute in commerce a product, or a flame-retarded part of a product, containing more than one-tenth of 1 percent of pentaBDE or octaBDE, by mass.”



Summary

- **PBDEs in consumer products are escaping into the environment – now everywhere**
- **PBDE levels rising rapidly in fish, other wildlife and people in North America**
- **Some folks have much higher levels than most for reasons unknown.**
 - Levels similar to levels associated with health effects
- **Penta- and Octa-PBDE banned in CA and the EU; not used in Japan.**
 - Renewed concern over Deca (new data on debromination by UV and biota)

